



# Plagiarism Checker X Originality Report

**Similarity Found: 17%**

Date: Friday, September 10, 2021

Statistics: 1238 words Plagiarized / 7138 Total words

Remarks: Low Plagiarism Detected - Your Document needs Optional Improvement.

---

Available online at <http://link.springer.com> Ocean Sci. J. (2017) 52(1):75 ? 89  
<http://dx.doi.org/10.1007/s12601-017-0010-y> pISSN 1738-5261 eISSN 2005-7172 Article  
Reproductive Biology of Blood Cockle *Anadara granosa* (Bivalvia: Arcidae) in the  
Northern Region of the Strait of Malacca Munawar Khalil 1\*, Zulfigar Yasin 2, and Tan  
Shau Hwai 2 1Department of Aquaculture, Universitas Malikussaleh, Aceh 24351,  
Indonesia 2Marine Sciences Laboratory, School of Biological Science, Universiti Sains  
Malaysia, Gelugor 11800, Malaysia Received 19 February 2016; Revised 4 August 2016;  
Accepted 31 October 2016 ? KSO, KIOST and Springer 2017 Abstract ? A study on the  
reproductive cycle of the blood cockle *Anadara granosa* (Bivalvia: Arcidae) was  
conducted at three different areas in the northern region of the Strait of Malacca. A total  
of 1,920 sa mples of adult A.

*granosa* (38–71 mm length) were collected from June 2009 until Septem ber 2010.  
Qualitative techniques (gonadal microscopic fresh smear test and histology analysis) as  
well as quantitative techniques (analysis of condition index and gonadal index) were  
used to predict monthly gonadal development stages of *A. granosa* . The gonadal index  
of *A. granosa* from Banda Aceh (Indonesia) (  $r = 0.469$ ,  $P > 0.05$ ) and Pulau Pinang  
(Malaysia) ( $r = 0.123$ ,  $P > 0.05$ ) did not show any corr elation to their condition index,  
whereas the gonadal index of *A.*

*granosa* from Lhokseumawe (Indonesia) ( $r = 0.609$ ,  $P < 0.05$ ) showed moderate positive  
correlation to the condition index. During the 16 month sampling period, four  
reproductive cycles were observed: each from three to six months. The process of  
releasing gametes is termed dribble spawning, and is the same in all populations. The  
principle component analysis (PCA) indicated that *A. granosa* reproduction was affected  
by interaction between internal physiological factors and indigenous environmental

factors. In all sampling areas, phytoplankton density played a key role in the reproductive cycle in *A. granosa*.

Information on the reproductive biology of this species is essential for species management and to improve the sustainability practices of the fisheries industry. These findings will provide basic information on the biology of the blood cockle *A. granosa* for stock management in the region. Key words ? blood cockle, reproductive cycle, gametogenesis, gonadal index, condition index 1. Introduction *Anadara granosa* is one of 7500 bivalve species in the family Arcidae, often called "blood arks" or "blood cockles" (Gosling 2003; Arapov et al. 2010).

Their common name refers to the hemoglobin and hemocyanin pigments in their blood and tissue cells, giving their blood a dark red color (Ruppert and Barnes 1994) which has allowed this species to live in oxygen-critical habitats (Broom 1985; Terwilliger and Terwilliger 1985; Cilenti et al. 2010). The species is indigenous to the intertidal mudflats of many Southeast Asian countries, particularly Indonesia, Malaysia and Thailand. *Anadara granosa* are mainly distributed in mangrove forests, muddy vegetation and mixed areas. The intertidal species *A. granosa* is known as a keystone species in mangrove habitats in several areas in the northern region of the Strait of Malacca.

This species has also been one of the most important fisheries commodities in Southeast Asia for many years (Borrero 1986; Broom 1985; Suwanjarat et al. 1990, 2009). The northern Strait of Malacca is an important nursery area for many intertidal organisms and a feeding area for migrating species. Being the most important species in terms of fisheries production, this cockle has become the target of an extensive culturing operation in West Malaysia (Broom 1983).

At the same time, harvesting of wild stock of cockles in Sumatra and Java (Indonesia) is at an all-time high to meet the demand for shellfish. In Malaysia, the annual production of blood cockles in 2009 exceeded metric 65,000 tonnes, which is valued at US \$36.60 million (DOF 2010). The main blood cockle production areas in Malaysia are concentrated in Kedah (Merbok), Pulau Pinang (Juru), Perak (Kuala Gula, Kuala Sangga-Matang, Kuala Trong, Sungai Jarum), Selangor (Kuala Selangor) and Johor (Muar).

In Indonesia, this species can be found in abundance on the \*Corresponding author. E-mail: khalil@unimal.ac.id 76 Khalil, M. et al. coast of West Sumatra, Central and South Java, East and West Kalimantan and other muddy bottoms in Sulawesi, Maluku and Papua (Khalil et al. 2009). The most recent data available on annual cockle production in Indonesia is from 2009 when it reached 47,437 metric tonnes, or equal to US\$ 23.72

million (DKP 2010).

The Northern Straits of Malacca is an important area for the harvesting and culture of the blood cockle *A. granosa* due to the suitability of the habitats for spawning and growth (Mirzaei and Hwai 2016). However, annual production statistics indicate a decrease in stocks in the last decade. This situation may be due to inadequate management of wild cockle populations. Fisheries management is needed to improve policies for the sustainability of the fisheries industry.

In depth information on reproductive cycles is necessary for predicting annual recruitment, as well as interpreting growth, mortality, and survival data in the marine culture of species (Shaw 1965; Manzi et al. 1985; Sbrenna and Campioni 1994). This data is lacking for the blood cockle *Anadara granosa* but is essential to optimize aquaculture of this species. This bivalve species can be managed more effectively after evaluating the regeneration capabilities of natural stocks and interpreting growth patterns.

Detailed and comprehensive information on gonadal development is also important for economic management of this species (Gribben et al. 2004; Peharda et al. 2006). This study aimed to investigate the seasonal gonadal cycle of the cockle *A. granosa* by using quantitative techniques (gonadal index and condition index) through gonadal fresh smear test and gonad histology (a qualitative technique) from specimens collected from the northern region of the Strait of Malacca. 2. Materials and Methods Collecting of samples A total of 120 samples of adult *A. granosa* were collected Fig. 1.

Sampling location Reproductive Biology of *Anadara granosa* 77 monthly from June 2009 till September 2010 from the natural habitat in Banda Aceh (532'34.67"N–9517'2.54"E), Lhokseumawe (0509'35.3"N–09708'29.4"E) in Aceh, Indonesia and Pulau Pinang (516'9.66"N–10023'27.37"E) in Malaysia (Fig. 1). The total number of specimens sampled was 1,920 and the adult cockle sizes ranged from 38–71 mm in length. The sampling area was characterized by muddy substrate which was surrounded by mangroves, no wave action and high salinity. The specimens were collected at a depth of 5–30 cm and salinity ranged from 10–33 ppt. Sampling activity in the field was conducted once a month over the specified time frame during low tides.

The live specimens were collected manually with the aid of a harrow, which was run through the muddy area to pull bivalves to the surface. After collecting, the specimens were stored in isotherm containers and immediately transported to the laboratory. The samples were cleaned to fully remove all fouling organisms and other adherences. Qualitative technique Gonadal microscopic fresh smear test A total of 40 specimens per sampling site were randomly allocated for the gonadal microscopic fresh smear test

each month. All the specimens were dissected with a dissecting needle and pipette.

The fresh smear procedure was adopted to observe the gonad content under a compound light microscope (magnification = 100 x) to analyze the stages of the gonadal development. The sex and gametogenesis stages were identified using image analysis, which included 4 stages: (+1) indeterminate, (+2) developing, (+3) developed and (+4) spawned (Rajagopal et al. 2006). Histology analysis A total of ten gonad specimens from each of the three sampling sites were allocated for histological analysis each month. Slides were prepared through the process of embedding paraffin wax into the tissue.

Haematoxyline and Eosin coloration were used for tissue coloring (Howard et al. 2004). The initial process requires dehydration of the specimen tissue. Dehydration was done through a series of steps of immersing the sample into varying concentrations of alcohol. The sample was embedded into a mold of wax next and kept in a refrigerator overnight before preparing it for HE coloration. The solutions used for histology included bouins, alcohol (50%, 70%, 80%, 90%, 95% and absolute alcohol), xylene, liquid wax, histosolve, HE solution and 1.5% ammonia.

A microtome was used to cut 5–7 µm thick tissue sections which were mounted on a glass microscope slide. The light compound microscope was used to analyze the gonad structure to recognize the sex and gametogenesis stages (divided into: (+1) indeterminate, (+2) developing, (+3) developed and (+4) spawned). Quantitative method Analysis of condition index (CI) The water displacement method was used to determine the condition index.

A total of 30 specimens (size range: 38– 71 mm in length ) from each sampling station were examined from June 2009 to September 2010. Each specimen was measured for the following: dry flesh weight, wet weight of shell in grams (g) and internal cavity volume (ml). Fresh cockle tissue including its shell was weighed using digital balances. The flesh was dried at 105°C for 72 hours to a constant weight. The volume of the shell internal cavity was calculated by means of subtracting the volume of the shell (ml) from the total wet volume (ml).

These data were used to calculate the condition index using the formula described by Lawrence and Scott (1982):  $\text{Condition index} = \frac{\text{dry flesh weight} \times 100}{\text{shell internal cavity volume (cm}^3)}$  Analysis of gonadal index (GI) Gonadal index was calculated based on the formula proposed by Gosling (2003) and Kim and Lee (2008):  $\text{Gonadal index} = \frac{\sum \text{individual from each stage level gonad stage}}{n \text{ total specimen for each sampling batch}}$  The gonadal index (GI) was calculated for each sampling month through the

gonadal microscopic fresh smear test and histological analysis to estimate the proportion of the gonadal stages (indeterminate, developing, developed and spawned).

The GI value was ranked to: 1 (all individuals' gonads in the samples were in spawned stage), 2 (all individuals' gonads in the samples were in indeterminate stage), 3 (all individuals' gonads in the samples were in developing stage) and 4 (all individuals' gonads in the samples were in developed stage). Statistical analysis Raw data was compiled and entered into Microsoft Office Excel 2011 (Macintosh version) for processing and analyzing of minimum and maximum values, averages, and the standard deviation as well as to generate graphs.

One-Way ANOVA statistical analysis and post hoc test were used to determine significance level ( $P < 0.05$  and  $P < 0.01$ ) in the values of each data cluster. Pearson correlation test was also utilized to determine and understand the relationship between differing variables (CI and GI). The principle component analysis (PCA) was used to analyze the correlation between parameters which were affected by reproductive activities in each sampling area.

These statistical analyses were conducted using SPSS (Statistical Package for Social Science) release 20.0 for Macintosh. 3. Results Gonadal structure of *Anadara granosa*  
Gonadal microscopic fresh smear analysis The description of gonad structure of *A. granosa* based on microscopic fresh smear analysis was categorized as shown below:  
Stage 1 (indeterminate). Male and female: Determination of sex cannot possibly be determined. Gonadal compound appeared to be empty and filled up only by network of connecting tissues. Unused residual of gametes can be found. Stage 2 (developing). Male: The gonadal compound turned cream in color.

Gametes have been very active and the testis was filled with spermatogonia and spermatid. Spermatozoa were also found in limited numbers and sometimes found in tailed form and actively swim. Female: The gonadal compound turned orange in color. Gametes in ovary have begun to appear, which are previtellogenic oogonia, oocytes and a limited number of oocytes vitellogenic. Oocytes were scattered and filled inside the follicle. Nuclei in oocytes vitellogenic have started forming and are clearly visible. Oocytes have uneven sizes. Stage 3 (developed). Male: The gonadal compound turned a more concentrated cream color as a result of highly condensed developed spermatozoa.

The spermatozoa have already developed their tail and are swimming actively. Sometimes, spermatids can still be found in small numbers. Female: Gonadal compound turned an intense, concentrated orange due to formation of highly condensed oocytes. Gametes were generally mature oocytes. Oocytes are in polyhedral form. The nuclei

within the oocytes have matured and grown larger in size. The yolks were found in most of the mature oocytes. Previtellogenic oocytes can still be found in small amounts. Stage 4 (spawned). Male: Gonadal compound reduced drastically. Spermatozoa have diminished. Unused residual spermatozoa can be found inside the lumen. Female: Gonadal compound turned bright orange due to the lowest concentration of oocytes.

Mature oocytes were found in small amounts, but these are expected to be residue or absorbed as phagocytes. Most of the oocytes had no shape and the nuclei appeared to have shrunk and disappeared. Gonadal histology analysis Stage 1 (indeterminate). Male and female: The stage is also called the dormant stage; the sexes cannot be distinguished. Undeveloped gonads' content during this stage only consisted of connecting tissues and a handful of residual gametes leftover from the previous spawned stage (stage 4) (Fig. 2). Stage 2 (developing).

Male: Gonad was gradually filled up with spermatogonia, spermatocyte, and a small quantity of spermatozoa. The average diameter of the follicles at this stage was  $117.77 \pm 19.58 \mu\text{m}$  in size (Fig. 3a). Female: Oocytes occur in a range of sizes and were generally not the same shape (irregular). Gonad was gradually filled up with oogonia as well as vitellogonia oocyte and vitellogenic oocytes, the nuclei have uneven shapes. The average diameter of the follicles at this stage was  $136.21 \pm 22.12 \mu\text{m}$ , whereas the average diameter of oocytes was  $24.81 \pm 6.19 \mu\text{m}$  in size (Fig. 4a). Fig. 2. Gonadal structure of *Anadara granosa* based on histology analysis at indeterminate stage.

FW: follicle wall; Lu: Lumen; EL: empty lumen; Ct: connective tissue Reproductive Biology of *Anadara granosa* 79 Stage 3 (developed). Male: Gonad was mainly dominated by spermatozoa content. Interfollicular space at this stage was seen to be experiencing constriction due to the growing follicle size. Spermatogonia were still found in limited number and typically found on the side wall of the follicle. The average diameter of the follicles was  $186.16 \pm 14.47 \mu\text{m}$  in size (Fig. 3b). Female: Gonad was characterized by the dominance of vitellogenic oocytes with a visibly large nucleus.

Lumen space was dominated by the polyhedral oocyte vitellogenic shape which was untouched or free from the follicle wall. The cytoplasm of mature oocytes had been filled by a number of yolk granules. The average diameter of the follicles was  $215.13 \pm 38.40 \mu\text{m}$  and oocytes were  $30.01 \pm 6.80 \mu\text{m}$  in size (Fig. 4b). Stage 4 (spawned). Male: Spermatozoa seemed to be reduced, as the follicle appeared almost empty. Spermatogonia were not found (Fig. 3c). Female: Residual oocytes were present. The follicles' wall seemed to be damaged and unfilled. Phagocytes were found around the residue oocytes (Fig. 4c).

Gonadal development cycle The gonad percentage (for each stage) was compared between the three sampling locations: Banda Aceh (Indonesia), Lhokseumawe (Indonesia) and Pulau Pinang (Malaysia). Figures 5a, 6a, 7a, as well as 5b, 6b and 7b depict the computation of gonad percentages per month for all the 4 phases discussed covering a span of 16 months, from June 2009 until September 2010, through gonadal microscopic fresh smear analysis Fig. 3. Gonadal structure of male *Anadara granosa* based on histology analysis: (a) Stage 2 (Developing); (b) Stage 3 (Developed); (c) Stage 4 (Spawned).

FW: follicle wall; Lu: lumen; Spz: spermatozoa; MS: mature spermatozoa; SD: sperm ductus; St: spermatid; DS: degenerative space; DSz: degenerative spermatozoa; FF: follicle fragment; EF: empty follicle; Ct: connective tissue Fig. 4. Gonadal structure of female *Anadara granosa* based on histology analysis: (a) Stage 2 (Developing); (b) Stage 3 (Developed); (c) Stage 4 (Spawned). FW: follicle wall; Lu: Lumen; EVO: early stage of vitellogenic oocyte; LVO: late stage of vitellogenic oocyte; MO: mature oocyte NI: nucleus; FF: follicle fragment; EF: empty follicle; RO: residual oocyte; IS: interfollicular space; YG: yolk granule 80 Khalil, M. et al. and gonadal histology analysis, respectively. Figures 5c, 6c and 7c, as well as 5d, 6d and 7d depict the monthly condition index (CI), and monthly gonadal index (GI), respectively, covering the same 16 months. Fig. 5.

*Anadara granosa* gonadal development pattern from Banda Aceh, Indonesia (June 2009–September 2010) Reproductive Biology of *Anadara granosa* 81 Environmental variable Monthly seasonal variations of environmental parameters in the three different sampling areas are reported in Table 1. During the study period, water temperature, salinity and phytoplankton density fluctuated significantly compared to other environmental parameters. The principle component analysis (PCA) was conducted to evaluate the comprehensive relationship between environmental factors and reproduction variables in the *A. granosa* populations in Banda Aceh, Lhokseumawe, and Pulau Pinang (Fig. 8). Fig. 6.

*Anadara granosa* gonadal development pattern from Lhokseumawe, Indonesia (June 2009–September 2010) 82 Khalil, M. et al. 4. Discussion Gonad development for *Anadara granosa* The recorded CI values for the samples indicated significant varying values every month for samples from the same sampling location as well as those from different sampling locations. The difference in the trend of CI value indicated the status of the population of blood cockles throughout the year. A high CI value implies the gonad has already reached maturity. However, CI is not always linearly correlated to its breeding pattern.

This can be shown from the comparison of the monthly CI vs GI values. The GI value is

an assumed indication of the breeding status. A sudden drop in GI value signifies the occurrence of spawning activities. From this analysis, there was no linear correlation between CI and GI values for samples from Banda Aceh and Penang. However, Fig. 7. *Anadara granosa* gonadal development pattern from Pulau Pinang, Indonesia (June 2009–September 2010) Reproductive Biology of *Anadara granosa* 83 Table 1.

Mean monthly seasonal environmental parameter at the sampling areas from June 2009 to September 2010 Environmental parameter Jun. 2009 Jul. 2009 Aug. 2009 Sep. 2009 Oct. 2009 Nov. 2009 Dec. 2009 Jan. 2010 Feb. 2010 Mar. 2010 Apr. 2010 May 2010 Jun. 2010 Jul. 2010 Aug. 2010 Sep. 2010 Average Temperature ( °C) Banda Aceh Minimum 26.32 24.27 24.29 25.43 26.90 25.00 25.39 25.79 22.96 22.84 25.47 21.87 25.90 22.82 25.27 26.00 24.78 Maximum 30.97 30.44 30.45 29.88 32.61 31.93 31.95 32.48 32.93 32.10 31.40 31.03 30.93 32.11 30.98 30.38 31.41 Lhokseumawe Minimum 28.82 28.82 26.81 28.07 27.71 27.03 28.45 28.06 28.96 28.48 28.60 28.23 27.95 28.23 27.23 27.40 28.05 Maximum 31.08 31.71 30.06 31.17 30.87 30.33 30.81 30.68 30.75 31.10 30.90 31.65 31.27 31.39 30.84 31.17 30.99 Pulau Pinang Minimum 27.23 27.52 26.65 27.10 25.87 27.93 26.05 23.68 25.61 26.74 26.37 26.29 26.67 26.74 26.65 27.37 26.53 Maximum 31.63 31.10 31.45 30.60 30.90 31.43 30.58 28.35 30.46 31.71 31.13 30.68 31.53 31.32 31.45 31.30 30.98 Salinity (ppt) Banda Aceh 32.27 31.35 29.98 27.47 30.06 27.20 26.45 29.68 31.50 31.16 29.30 30.71 30.85 31.29 30.45 28.27 29.87 Lhokseumawe 31.00 30.97 31.16 31.20 31.03 29.07 30.94 31.16 31.46 31.84 30.57 31.26 30.90 30.81 30.65 31.27 30.95 Pulau Pinang 29.33 28.52 26.39 26.87 27.35 26.13 25.23 28.06 29.00 28.94 28.67 26.48 29.70 29.32 31.06 30.40 28.22 pH Banda Aceh 7.65 8.02 8.03 8.17 7.80 8.02 7.91 8.17 8.08 8.06 8.02 7.97 7.74 8.27 7.94 8.23 8.01 Lhokseumawe 8.13 7.88 8.04 8.06 8.17 8.13 7.98 8.21 7.89 8.18 7.91 7.84 8.08 7.93 7.99 8.13 8.03 Pulau Pinang 8.02 7.49 7.85 8.14 8.04 8.07 7.79 8.02 7.95 7.86 8.08 8.21 7.91 7.86 7.33 7.54 7.89 Dissolved oxygen (mg/L) Banda Aceh 6.53 6.81 6.96 6.05 5.97 6.12 6.05 5.95 6.10 5.84 6.23 5.86 6.68 5.84 6.32 5.47 6.17 Lhokseumawe 6.01 6.38 6.47 6.14 6.28 6.04 6.07 6.10 6.97 6.02 6.17 5.98 6.02 5.89 6.28 6.13 6.18 Pulau Pinang 7.20 5.20 5.20 4.90 5.13 5.21 5.09 5.29 5.20 5.64 5.87 5.39 7.67 5.64 6.29 5.87 5.67 Turbidity (NTU) Banda Aceh 17.40 29.30 8.61 9.02 10.86 9.12 19.16 14.09 16.03 10.27 18.98 13.83 10.27 34.29 9.74 10.48 15.09 Lhokseumawe 43.20 30.50 36.50 66.90 31.60 15.18 103.00 93.67 29.13 37.30 64.92 38.95 35.30 49.98 46.90 98.30 51.33 Pulau Pinang 29.30 17.36 15.11 13.09 17.27 74.30 57.80 77.10 109.67 107.00 98.00 76.00 103.40 107.00 76.65 93.12 67.01 Orthophosphate (mg/L) Banda Aceh 0.05 0.03 0.04 0.13 0.03 0.02 0.00 0.40 0.00 0.07 0.13 0.07 0.03 0.09 0.08 0.53 0.11 Lhokseumawe 0.05 0.01 0.02 0.07 0.01 0.01 0.01 0.70 0.00 0.04 0.06 0.04 0.04 0.08 0.07 0.01 0.08 Pulau Pinang 0.10 0.05 0.08 0.06 0.03 0.01 1.00 0.01 0.16 0.13 0.09 0.08 0.52 0.13 0.15 0.81 0.21

84 Khalil, M. et al. Table 1. Continued Environmental parameter Jun. 2009 Jul. 2009 Aug. 2009 Sep. 2009 Oct. 2009 Nov.



2009 Dec. 2009 Jan. 2010 Feb. 2010 Mar. 2010 Apr. 2010 May 2010 Jun. 2010 Jul. 2010 Aug. 2010 Sep. 2010 Average Nitrate (mg/L) Banda Aceh 0.71 0.03 0.11 0.73 0.75 0.01 0.04 0.05 0.05 0.09 0.23 0.63 0.18 0.05 0.53 0.65 0.30 Lhokseumawe 0.68 0.14 0.03 0.03 0.20 0.01 0.00 0.03 0.01 0.10 0.77 0.58 0.10 0.17 0.07 0.98 0.24 Pulau Pinang 0.80 0.02 1.30 0.73 0.03 2.02 0.03 1.02 0.11 0.14 0.61 0.42 0.64 0.14 0.65 1.76 0.65 Nitrite (mg/L) Banda Aceh 0.05 0.02 0.02 0.03 0.75 0.01 0.03 0.03 0.03 0.03 0.05 0.05 0.05 0.03 0.04 0.05 0.08 Lhokseumawe 0.03 0.03 0.04 0.04 0.05 0.03 0.44 0.03 0.43 0.02 0.08 0.07 0.02 0.08 0.03 0.04 0.09 Pulau Pinang 0.03 0.03 0.04 0.06 1.09 1.68 0.18 1.00 0.13 0.12 0.09 0.03 0.14 0.12 0.39 0.87 0.37 Ammonia (mg/L) Banda Aceh 0.87 0.20 0.16 0.15 0.11 0.68 0.06 0.13 0.08 0.09 0.19 0.79 0.18 0.06 0.59 0.13 0.28 Lhokseumawe 0.19 0.17 0.25 0.25 0.14 0.35 0.23 0.30 0.19 0.27 0.39 0.49 0.27 0.21 0.43 0.24 0.27 Pulau Pinang 0.24 0.18 0.14 0.25 0.42 0.11 0.61 0.15 0.68 0.54 0.65 0.98 0.65 0.54 0.82 0.65 0.48 Phytoplankton density (cell/L) Banda Aceh 1831.67 1446.67 851.67 1178.33 630.00 991.67 1201.67 385.00 1773.33 1388.33 1516.67 1785.00 1738.33 1283.33 1341.67 1108.33 1278.23 Lhokseumawe 1656.67 1365.00 711.67 2601.67 3010.00 1435.00 4001.67 1365.00 2986.67 2415.00 2333.33 2298.33 2415.00 1050.00 1003.33 2310.00 2059.90 Pulau Pinang 4340.00 4001.67 1470.00 11713.33 4340.00 4281.67 2636.67 4561.67 4235.00 5751.67 5693.33 6090.00 5728.33 5751.65 2905.00 7910.00 50 88.12 Reproductive Biology of *Anadara granosa* 85 a linear correlation between these values was noted for samples from Lhokseumawe. These were tested with the Pearson correlation test, which indicated that CI values for samples from Banda Aceh ( $r = 0.469$  at  $P > 0.05$ ) and Penang ( $r = 0.123$  at  $P > 0.05$ ) has no significant correlation to their respective GI, but there is a mild correlation for samples from Lhokseumawe ( $r = 0.609$  at  $P < 0.05$ ).

A negative correlation has also been reported in studies of other bivalve species. Herrmann et al. (2009) reported a negative correlation between CI and gametogenesis cycle for *Amarilladesma mactroides* (Reeve, 1854). Mladineo et al. (2007) also reported zero correlation between CI and GI for the bivalve *Modiolus barbatus* (Linnaeus, 1758). The same applies to *Mercenaria mercenaria* (Linnaeus, 1758) from the Gulf of Narragansett in the United States, as reported by Marroquin-Mora and Rice (2008).

The GI values obtained throughout the year indicate high diversity in reproductive patterns among the three sampling locations. This is expected due to the differences in habitat conditions as well as breeding seasons. Blood cockles from all three sampling locations showed a rapid transition from gonad development to maturation phase. GI analysis shows that spawning activity happened every month throughout the year with varying intensity. The GI value increases during gametogenesis and decreases after spawning. The fast-paced transition could be a strategy for the blood cockles to increase the amount of gamete released whilst favorable environmental conditions are present.

This behavior is characteristic of the reproduction of invertebrates in tropical regions. Species have been shown to adopt opportunistic strategies to develop the gonadal matter from energy available from food rather than from energy stored inside somatic parts (Cárdenas and Aranda 2000). Freitas et al. (2010) found that *Anadara notabilis* exhibits a continuous reproductive cycle throughout the year and that particulate organic matter, temperature and food availability were regulating factors of the reproduction of *A. notabilis*.

This study of blood cockles' GI shows that it has a breeding cycle lasting an average of 3~6 months across the three sampling locations (Banda Aceh, Lhokseumawe and Pulau Pinang). During the 16 month sampling period, four reproductive cycles were observed. For the *A. granosa* population from Banda Aceh (Indonesia), cycle I occurred from June to October 2009, cycle II from November 2009 to January 2010, cycle III from February to April 2010, and cycle IV from April to September 2010. In Lhokseumawe (Indonesia), cycle I started from June to August 2009, cycle II from September 2009 to January 2010, cycle III from February to June 2010 and cycle IV from July to September 2010. For the *A.*

*granosa* population in Pulau Pinang (Malaysia), cycle I started from June to October 2009, cycle II from November 2009 to February Fig. 8. Principle component analysis (PCA) plot for the reproductive factor component on *Anadara granosa* population. (a) Banda Aceh, Indonesia; (b) Lhokseumawe, Indonesia; (c) Pulau Pinang, Malaysia 86 Khalil, M. et al. 2010, cycle III from February 2010 to April 2010 and cycle IV from April 2010 to September 2010. All three populations started the first cycle around June 2009 and ended the fourth cycle also around the same time, September 2010. The population from Lhokseumawe (Indonesia) showed a tendency to spawn faster compared to the other two populations.

However, during the third cycle, populations from Banda Aceh (Indonesia) and Penang (Malaysia) exhibited a more rapid and shorter cycle lasting approximately 2~3 months, compared to Lhokseumawe (Indonesia) which took about 5 months. Breeding pattern of *Anadara granosa* Generally, the bivalve breeding process is characterized by a continual and seasonal pattern (Ceballos-Vazquez et al. 2000), and is iteroparous in nature, continually and repeatedly breeding throughout its entire life span (Dame 1996). Bivalves give birth to their young by means of gametogenesis. This process is then followed by the release of one or several gametes.

The process of rearranging the empty gonad with new gametes for the next cycle always is the signal for the beginning of a new breeding cycle (Gosling 2003). Variation in the

breeding trend amongst cockle populations of different geographical locations makes it difficult to determine a pattern of gonad development. A well balanced distribution of males to females in blood cockles is supported by the sex ratio analysis done in this study. Gonad development and spawning period was determined to be parallel between the two opposing sexes, a scenario known as synchrony.

According to Levitan (1993), synchrony in gonad development of bivalves is crucial to increase the possibility of effective mating. Extended spawning durations from one to two months is a common breeding strategy for bivalve species. Such a strategy is essential to maintain the cockle population over time within its habitat. Generally, sporadic gamete mating will happen concurrently under suitable surrounding conditions. Blood cockles for all three sampling locations, and in general, exhibit a tendency to be characterized as bivalve brachidictics, which means they are capable of undergoing a continual breeding cycle throughout the year, with varying spawning intensity every month. Pathansali (1966), Narasimham (1988) and Broom (1983) reported that *A. granosa* in Peninsular Malaysia and India has a spawning season throughout the year with no apparent seasonal pattern.

In comparison, the spawning season of Archidae (genus *Anadara*) is presented in Table 2. The information on the reproductive cycle of *A. granosa* provided by this study is crucial for initiating its commercial aquaculture value, as well as for the sustainable management of wild stocks. In the future, data on spawning periodicity might be used to identify trochophore or veliger larvae in wild habitats and for seed collection activities.

When bivalve culture production depends on natural seed supply, the timing of seed collection is critical since the potential brood stock is Table 2. Comparison of spawning period with the highest intensity of releasing gamete in genus *Anadara*

Species	Location	Spawning period	Sources
<i>A. granosa</i>	Banda Aceh, Indonesia	September and October	Present study
<i>A. granosa</i>	Lhokseumawe, Indonesia	Jun and September	Present study
<i>A. granosa</i>	Pulau Pinang, Malaysia	October, February to March	Present study
<i>A. granosa</i>	Perak	December	Pathansali (1966)
<i>A. granosa</i>	Phuket, Thailand	October to November	Boonruang and Janekarn (1983)
<i>A. granosa</i>	Pulau Pinang, Malaysia	August to September	Broom (1983)
<i>A. granosa</i>	Selangor, Malaysia	September and November	Broom (1983)
<i>A. granosa</i>	West coast, Thailand	August and November	Suwanjawat and Pannrong (1990)
<i>A. granosa</i>	Pattani bay, Thailand	September, December and July	Suwanjarat et al. (2009)
<i>A. trapezia</i>	Sydney, Australia	end of summer	Sullivan (1961)
<i>A. broughtoni</i>	Japan	Jun to August	Kanno and Kikuchi (1962)
<i>A. senilis</i>	Nigeria coast	October and November	Yoloye (1974)

broughtoni Great Bay, Japan July to September Dzyuba and Maslennikova (1982) A.

trigonopsis New Zealand winter and summer Booth (1983) A. trapezia Sydney water, Australia winter Hadfield and Anderson (1988) A. descripta Sydney water, Australia winter and autumn Hadfield and Anderson (1988) A. inaequalis Black sea, Turkey Mei and November Sahin et al. (2006) Anadara/Scapharca kagoshimensis Ariake bay, Japan autumn to early of summer Yurimoto et al. (2008) A. notabilis Northeastern of Venezuela Jun and October Freitas et al. (2010) Reproductive Biology of Anadara granosa 87 primed for a short period of time. Information presented here indicates that quantitative methods (condition index and gonadal index) are a precise indicator in A. granosa brood stock.

Factors that affected reproduction cycle of Anadara granosa in the northern region of the Strait of Malacca Gametogenesis is affected by the change and interaction of exogenous (temperature, salinity, light, food), and endogenous factors (nervous system, hormones) that could determine the reproductive strategy of bivalve species (Ram et al. 1996; Utting and Millican 1997; Louro et al. 2003; Barber and Blake 2006; Magnesen and Christophersen 2008). The principle component analysis (PCA) has shown that the principle components which affected the reproductive cycle of the A.

granosa population in Banda Aceh were gonadal index, condition index, phytoplankton density, orthophosphate, salinity, and water temperature. The principle component analysis for Lhokseumawe showed that there were five variables affecting A. granosa reproduction, namely interaction among gonadal index, condition index, phytoplankton density, ammonia, and pH. The reproduction of the A. granosa population in Pulau Pinang also revealed the complex interaction of seven principle variables, namely interaction between gonadal index and environmental factors such as salinity, nitrite, ammonia, phytoplankton density, turbidity and dissolved oxygen. Principle component analysis indicated that the environment parameters modifying the reproduction of A. granosa populations were diverse and complex. This analysis also indicated that reproduction of A.

granosa populations is significantly affected by the interaction of local environment parameters. For example, water temperature was found to be modestly interacting with and affecting components of reproduction in A. granosa in Banda Aceh. Dissolved oxygen, nitrite, ammonia, and turbidity variables were only found to be specifically interacting and affecting the components of reproduction in the A. granosa population in Pulau Pinang, however these variables were not the factors affecting reproduction in the A. granosa population in Banda Aceh and Lhokseumawe.

Reproductive physiological factors such as gonadal index and condition index were shown to be affected only by the interaction of several water environment parameters that are dependent on adaptation level (Gillmor 1982; Beninger and Le Pennec 1997). One of the environmental factors that is known to be strongly correlated with *A. granosa* reproduction is phytoplankton density. This variable is known to interact with and affect gonadal index and condition index as determined by gonadal development stage in the three *A. granosa* populations.

Lodeiros and Himmelman (1999) conducted a statistical analysis, namely multiple regression analysis, to see the relationship between environmental factors and reproduction of the bivalve *Lima scabra*. The conclusion of that study found that phytoplankton density was the only primary factor positively correlated to the reproduction of *L. scabra*. Phytoplankton density is the principle factor influencing the reproduction of bivalves (Wacker and von Elert 2003; Villalejo-Fuerte et al. 2005; Kang et al. 2006; Liu et al. 2006; Hernández-Olalde et al. 2007; Calderon-Aguilera et al. 2010).

Phytoplankton are also known to be the main source of diet to anadromous animals (Kasigwa and Mahika 1991). Gonadal maturation and the fertilization activities of *A. granosa* that correspond with the high level of phytoplankton density is a strategy to increase planktotrophic larval autonomy by increasing the larvae growth rate. The duration of the planktonic phase can be reduced through optimal utilization of the food source (phytoplankton).

Himmelman (1975) showed that a high content of phytoplankton in the aquatic environment will stimulate the reproductive period of invertebrate organisms, particularly species that have pelagic larvae. Ram et al. (1992) found that phytoplankton release a type of chemical substance that could stimulate the nervous system of bivalves to make them release gametes. Acknowledgements This project was supported by Universiti Sains Malaysia- Postgraduate Research Grant Scheme.

The authors would like to thank to Marine Sciences Laboratory Universiti Sains Malaysia, Ministry of Research Technology and Higher Education of the Republic of Indonesia, Universitas Malikussaleh Indonesia, Malaysia Quarantine and Inspection Services (MAQIS) Malaysia and Indonesia Fisheries Quarantine Service for their continuous support in making this project a success. References Arapov J, Ezgeta-Balic D, Peharda M, Ninčević Gladan Ž (2010) Bivalve feeding—how and what they eat? *Croat J Fish* 68(3): 105–116 Barber BJ, Blake NJ (2006) Reproductive physiology.

In: Shumway SE, Parsons GJ (eds) *Scallops: biology, ecology and aquaculture*, 88 Khalil, M. et al. Vol.35. Elsevier Sciences, New York, pp 357–416 Beninger PG, Le Pennec M

(1997) Reproductive characteristics of a primitive bivalve from a deep-sea reducing environment: giant gametes and their significance in *Acharax alinae* (Cryptodonta: Solemyidae). *Mar Ecol-Prog Ser* 157:195–206 Boonruang P, Janekarn V (1983) Distribution, density, biomass, and population bionomics of *Anadara granosa* (L.) in relation to environmental factors at Sapum Bay on the east coast of Phuket Island.

Thai Fish Gaz 36:461–468 Booth JD (1983) Studies on twelve common bivalve larvae, with notes on bivalve spawning seasons in New Zealand. *New Zeal J Mar Fresh* 17(3):231–265 Borrero FJ (1986) The collection of early juveniles of *Anadara* spp. as a potential source of seed for culturing mangrove cockles on the Pacific coast of Colombia. *Aquaculture* 59(1):61–69. doi:10.1016/0044-8486(86)90079-7 Broom M (1983) Mortality and production in natural, artificially-seeded and experimental populations of *Anadara granosa* (Bivalvia: Arcidae). *Oecologia* 58(3):389–397. doi:10.1007/BF00385241 Broom M (1985) The biology and culture of marine bivalve mollusc of the genus *Anadara*.

International Center for Living Aquatic Resources Management, Manila, ICLARM Studies and Reviews 12, 37 p Calderon-Aguilera LE, Aragón-Noriega EA, Reyes-Bonilla H, Paniagua-Chavez CG, Romo-Curiel AE, Moreno-Rivera VM (2010) Reproduction of the Cortes geoduck *Panopea globosa* (Bivalvia: Hiatellidae) and its relationship with temperature and ocean productivity. *J Shellfish Res* 29 (1):135–141 Cárdenas EB, Aranda DA (2000) A review of reproductive patterns of bivalve mollusks from Mexico. *B Mar Sci* 66(1):13–27 Ceballos-Vazquez BP, Arellano-Martinez M, Garcia-Dominguez F, Villalejo-Fuerte M (2000) Reproductive cycle of the rugose pen shell, *Pinna rugosa* Sowerby, 1835 (Mollusca: bivalvia) from Bahia Concepcion, Gulf of California and its relation to temperature and photoperiod. *J Shellfish Res* 19(1):95–99 Cilenti L, Scirocco T, Specchiulli A, Florio M, Renzi M, Breber P (2010) Population structure and spatial distribution of *Loripes lacteus* (Linnaeus, 1758) in Varano lagoon, SE Italy.

*Transit Water Bull* 2(4):63 ? 70. doi:10.1285/i1825229Xv2n4p63 Dame RF (1996) Ecology of marine bivalves: An ecosystem approach. CRC Press, Boca Raton, 272 p Dzyuba SM, Maslennikova LA (1982) Reproductive cycle of bivalve mollusks *Anadara broughtoni* in the southern part of Peter the Great Bay (Sea of Japan). *Soy J Mar Biol* 3:148–155 Freites L, Montero L, Arrieche D, Babarro JMF, Saucedo PE, Cordova C, García N (2010) Influence of environmental factors on the reproductive cycle of the eared ark *Anadara notabilis* (Röding, 1798) in northeastern Venezuela.

*J Shellfish Res* 29(1): 69–75. doi:10.2983/035.029.0128 Gillmor RB (1982) Assessment of intertidal growth and capacity adaptations in suspension-feeding bivalves. *Mar Biol* 68(3): 277–286 Gosling E (2003) Bivalve molluscs: biology, ecology and culture .

Blackwell, London, 456 p Gribben PE, Helson J, Jeffs AG (2004) Reproductive cycle of the New Zealand geoduck, *Panopea zelandica*, in two north island populations. *Veliger* 47(1):53–65 Hadfield AJ, Anderson DT (1988) Reproductive cycles of the bivalve molluscs *Anadara trapezia* (Deshayes), *Venerupis crenata* Lamarck and *Anomia descripta* Iredale in the Sydney region.

Mar Freshwater Res 39(5):649–660 Hernández-Olalde LNA, García-Domínguez FICO, Arellano- Martínez MAL, Ceballos-Vázquez BAP (2007) Reproductive cycle of the Pearl oyster *Pteria sterna* (Pteriidae) in the Ojo de Liebre lagoon, BCS, Mexico. *J Shellfish Res* 26 (2):543–548 Herrmann M, Alfaya JEF, Lepore ML, Penchaszadeh PE, Laudien J (2009) Reproductive cycle and gonad development of the Northern Argentinean *Mesodesma mactroides* (Bivalvia: Mesodesmatidae). *Helgoland Mar Res* 63(3):207–218. doi:10.1007/s10152-009- 0150-2 Himmelman JH (1975) Phytoplankton as a stimulus for spawning in three marine invertebrates.

J Exp Mar Biol Ecol 20(2):199– 214 Howard DW, Lewis EJ, Keller BJ , Smith CS (2004) *Histological techniques for marine bivalve mollusks and crustaceans*, 2nd edition. NOAA/National Centers for Coastal Ocean Science, Oxford, NOAA Technical Memorandum NOS NCCOS 5, 218 p DOF (2010) Perangkaan tahunan perikanan 2009 (Annual fisheries statistics 2009). Jabatan Perikanan Malaysia (DOF), Putra Jaya, Annual Fisheries Statistics Book, 26 p Kang CK, Lee YW, Choy EJ, Shin JK, Seo IS, Hong JS (2006) Microphytobenthos seasonality determines growth and reproduction in intertidal bivalves.

Mar Ecol-Prog Ser 315:113–127 Kanno H, Kikuchi S (1962) On the rearing of *Anadara broughtonii* (Schrenk) and *Haliotis discus hannai* Ino. *Bull Mar Biol Stat Asamushi* 11(2):71–76 Kasigwa PF, Mahika CG (1991) The diet of the edible cockle *Anadara antiquata* L. (Bivalvia, Arcidae) in Dar es Salaam, Tanzania, during the northeast monsoons. *Hydrobiologia* 209(1):7–12 DKP (2010) Buku statistik perikanan Indonesia 2010 (Indonesia annual fisheries statistics book 2010).

Ministry of Marine Affairs and Fisheries Republic Indonesia (DKP), Jakarta, Annual Fisheries Statistics Book 11, 182 p Khalil M, Yasin Z, Aileen TSW (2009) The current status of *Anadara granosa* in the east coast Sumatera island, Indonesia. In: *Proceedings of the Nagisa JSPS conferences, United Nation University, Tokyo, 15–16 Dec 2009* Kim TH, Lee KY (2008) Reproductive cycle and first sexual maturity of *Sinonovacula constricta* (Lamarck, 1818) (Bivalvia: Pharidae) in Western Korea. *Korean J Malacol* 24(2):97–104 Lawrence DR, Scott GI (1982) The determination and use of Reproductive Biology of *Anadara granosa* 89 condition index of oysters. *Estuaries* 5(1):23–27. doi:10.2307/1352213 Levitan DR (1993) The importance of sperm limitation to the evolution of egg

size in marine invertebrates.

Am Nat 141(4):517–536 Liu B, Dong B, Tang B, Zhang T, Xiang J (2006) Effect of stocking density on growth, settlement and survival of clam larvae, *Meretrix meretrix*.

Aquaculture 258(1):344–349 Lodeiros CJ, Himmelman JH (1999) Reproductive cycle of the bivalve *Lima scabra* (Pterioidea: Limidae) and its association with environmental conditions. Rev Biol Trop 47(3):411–418 Louro A, De la Roche JP, Campos MJ, Roman G (2003) Hatchery rearing of the black scallop, *Chlamys varia* (L.).

J Shellfish Res 22(1):95–100 Magnesen T, Christophersen G (2008) Reproductive cycle and conditioning of translocated scallops *Pecten maximus* from five broodstock populations in Norway. Aquaculture 285(1):109–116 Manzi JJ, Bobo MY, Burrell JR VG (1985) Gametogenesis in a population of the hard cockle, *Mercenaria mercenaria* (Linnaeus), in North Santee Bay, South Carolina. Veliger 28(2):186–194 Marroquin-Mora DC, Rice MA (2008) Gonadal cycle of northern quahogs, *Mercenaria mercenaria* (Linnaeus, 1758), from fished and non-fished subpopulations in Narragansett Bay.

J Shellfish Res 27(4):643–652 Mirzaei MR, Hwai AT S (2016) Assessing cockle shells (*Anadara granosa*) for reconstruction subdaily environmental parameters: implication for paleoclimate studies. Hist Biol 28(7):896–906. doi:10.1080/08912963.2015.1052806 Mladineo I, Peharda M, Orhanovi S, Bolotin J, Pavela-Vran IM, Treursi B (2007) The reproductive cycle, condition index and biochemical composition of the horse-bearded mussel *Modiolus barbatus*. Helgoland Mar Res 61(3):183–192. doi:10.1007/s10152-007-0065-8 Narasimham KA (1988) Biology of the blood clam *Anadara granosa* (Linnaeus) in Kakinada Bay. J Mar Biol Assoc India 30:137–150 Pathansali D (1966) Notes on the biology of the cockle *Anadara granosa* L.

In: Proceedings of the Indo-Pacific Fisheries Council 11, pp 84–98 Peharda M, Mladineo I, Bolotin J, Kekez L, Skaramuca B (2006) The reproductive cycle and potential protandric development of the Noah's Ark shell, *Arca noae* L.: implications for aquaculture. Aquaculture 252(2–4):317–327. doi:10.1016/j.aquaculture.2005.07.007 Rajagopal S, Venugopalan VP, Van der Velde G, Jenner HA (2006) Greening of the coasts: A review of the *Perna viridis* success story. Aquat Ecol 40(3):273–297. doi:10.1007/s10452-006-9032-8 Ram JL, Fong P, Croll RP, Nichols SJ, Wall D (1992) The zebra mussel (*Dreissena polymorpha*), a new pest in North America: reproductive mechanisms as possible targets of control strategies.

Invertebr Reprod Dev 22(1–3):77–86 Ram JL, Fong PP, Garton DW (1996) Physiological aspects of zebra mussel reproduction: maturation, spawning, and fertilization. Am Zool 36(3):326–338 Ruppert EE, Barnes RD (1994) Invertebrate zoology (6th edition).



Saunders College Publishing, Fort Worth, 1100 p Sahin C, Düzgünes E, Okumus I (2006) Seasonal variations in condition index and gonadal development of the introduced blood cockle *Anadara inaequalis* (Bruguiere, 1789) in the Southeastern Black Sea coast. *Turk J Fish Aquat Sc* 6(2):155– 163 Sbrenna G, Campioni D (1994) Gametogenic and spawning patterns of Manila cockles *Tapes philippinarum* (Bivalvia: Veneroida) in two lagoons of the River Po Delta, Italy.

*J Shellfish Res* 13(1):37–46 Shaw WN (1965) Seasonal setting patterns of five species of bivalves in the Tred Avon River, Maryland. *Chesapeake Sci* 6(1):33–37 Sullivan GE (1961) Functional morphology, micro-anatomy, and histology of the 'Sydney cockle' *Anadara trapezia* (Deshayes) (Lamellibranchia: arcidae). *Aust J Zool* 9(2):219–257 Suwanjart J, Parnrong S (1990) Reproductive cycles of *Anadara granosa* L. in Jebilung Satun Province.

Songklanakarin *J Sci Technol* 12(4):341–351 Suwanjart J, Pituksalee C, Thongchai S (2009) Reproductive cycle of *Anadara granosa* at Pattani Bay and its relationship with metal concentrations in the sediments. *Songklanakarin J Sci Technol* 31(5):471–479 Terwilliger RC, Terwilliger NB (1985) Molluscan hemoglobins. *Comp Biochem Phys B* 81(2):255–261. doi:10.1016/0305- 0491(85)90310-4 Utting SD, Millican PF (1997) Techniques for the hatchery conditioning of bivalve broodstocks and the subsequent effect on egg quality and larval viability.

*Aquaculture* 155(1):45–54 Villalejo-Fuerte M, Muneton-Gomez MS, Garate -Lizarraga I, Garcia-Dominguez F (2005) Gut content, phytoplankton abundance and reproductive season in the black oyster (*Hyotissa hyotis*, Linné, 1758) at Isla Espiritu Santo, Gulf of California. *J Shellfish Res* 24(1):185–190 Wacker A, von Elert E (2003) Food quality controls reproduction of the zebra mussel (*Dreissena polymorpha*). *Oecologia* 135(3): 332–338 Yoloye V (1974) The sexual phases of the West African bloody cockle *Anadara senilis* (L.).

*J Mollus Stud* 41 (1):25–27 Yurimoto T, Mori Y, Ito S, Maeno Y (2008) Reproductive cycle of the sub crenated ark shell *Scapharca kagoshimensis* (Tokunaga, 1906) in Ariake Bay, Japan. *J Shellfish Res* 27(5):1101–1108

#### INTERNET SOURCES:

-----  
<1% - [www.science.gov](http://www.science.gov) > topicpages > e

14% - [link.springer.com](http://link.springer.com) > article > 10

<1% - [www.researchgate.net](http://www.researchgate.net) > publication > 278814507

<1% - [www.researchgate.net](http://www.researchgate.net) > publication > 232684199

1% - [singapore.biodiversity.online](http://singapore.biodiversity.online) > taxo4254 > mainSpace

<1% - [www.sciencedirect.com](http://www.sciencedirect.com) › abs › pii

<1% - [www.jstor.org](http://www.jstor.org) › stable › pdf

<1% - [hmr.biomedcentral.com](http://hmr.biomedcentral.com) › 10 › s10152/009/0161-z

<1% - [www.tandfonline.com](http://www.tandfonline.com) › doi › figure

<1% - [www.researchgate.net](http://www.researchgate.net) › profile › Ismael-Garate