



Plagiarism Checker X Originality Report

Similarity Found: 15%

Date: Friday, May 20, 2022

Statistics: 722 words Plagiarized / 4752 Total words

Remarks: Low Plagiarism Detected - Your Document needs Optional Improvement.

N U S A N T A R A B I O S C I E N C E ISSN: 2087-3948 Vol. 14, No. 1, pp. 34-39 E-ISSN: 2087-3956 May 2022 DOI: 10.13057/nusbiosci/n140104 Agronomic diversity of several soybean putative mutant lines resulting from gamma-rays irradiation in M6 generation NILAHAYATI 1?, NAZIMAH1, RD. SELVY HANDAYANI1, JAKA SYAHPUTRA2, MUHAMMAD RIZKY2 1Department of Agroecotechnology, Faculty of Agriculture, Universitas Malikussaleh. Jl. Cot Tgk Ni Reuleut Timu, North Aceh 24355, Aceh, Indonesia. Tel./fax. +62-645-41373, email: nilahayati@unimal.ac.id 2Student Department of Agroecotechnology, Faculty of Agriculture, Universitas Malikussaleh. Jl. Cot Tgk Ni Reuleut Timu, North Aceh 24355, Aceh, Indonesia Manuscript received: 24 December 2021. Revision accepted: 13 February 2022. Abstract.

Nilahayati, Nazimah, Handayani RDS, Syahputra J, Rizky M. 2022. Agronomic diversity of several soybean putative mutant lines resulting from gamma-rays irradiation in M6 generation. Nusantara Bioscience 14: 34-39. Soybean is one of the foremost commodities for Indonesian people. Therefore, increasing domestic production absolutely must continue to be pursued. One way that can be done is to use the assembly technology of new superior varieties with better and adaptive properties.

This study aims to determine the agronomic diversity and yield of soybean putative mutant lines resulting from gamma irradiation in M6 generation. The research was conducted in Paloh Lada Village, Dewantara Subdistrict, Aceh Utara District, Indonesia, from July 2021 to October 2021. The tested genotypes consisted of 8 genotypes, namely Kipas Putih variety (parent variety), Anjasmoro variety (comparison variety), M.1.1.3, M.5.2.1, M.5.2.3, M.1.1.8, M.1.1.9 and M.1.1.17 mutant lines.

The results showed that the agronomic diversity of the tested mutants was different

compared with the parents and comparison variety. We obtained **early maturity and high yielding mutant** in M7 generation. The mutant lines with high yield and equivalent to its parents was M.1.1.8 with a production of 4.37 tha⁻¹. Mutant lines with early maturity were M.1.1.9, M.5.2.1 and M.1.1.3 lines with a harvesting age of 83 days after planting (DAP). Keywords: Gamma-ray irradiation, genotype, M6 generation, putative mutant lines INTRODUCTION Soybean is an important plant for Indonesian people (Lestari et al. 2017; Toyip et al. 2019).

Increasing national soybean production is necessary to maintain food sovereignty (Harsono and Pratiwi 2017). Various efforts can be made to increase domestic soybean production by expanding the planting area, improving cultivation techniques, and using superior varieties. Assembling superior varieties can be done by plant breeding techniques to obtain genetic diversity. Various methods of plant breeding, both conventional and biotechnology are currently available.

One of the very popular breeding techniques is mutation breeding with gamma-ray irradiation. Mutation induction with gamma-rays is commonly used to obtain genetic diversity in cultivated plants, especially soybeans. Several researchers have previously used gamma-rays on various soybean genotypes, including Sohag, BARI Soybean-5, Bangladesh Soybean-4, and BAU-S/64 for genetic improvement of yield attributes (Malek et al. 2014), soybean line VX04-6828 for better agronomic performance and seeds composition (Nobre et al. 2019), Anjasari soybean for high seed yield and large seed size (Mawarni et al.

2019; Yoel and Rachmadi 2020), Korean cultivars Danbaek and Daepung for highest linolenic acid contents (Hong et al. 2019), Kipas Putih soybeans for **early maturity and high** yields (Nilahayati et al. 2016; Nilahayati et al. 2019), Denna 2 variety for shade-tolerant (Harsanti et al. 2020) and induction **of genetic diversity of soybean** genotypes PGRI-15, KY EXOTIC, NARC-, AJMERI-2, AJMER-I (Hussain et al. 2020). It is possible to assemble local soybean varieties to obtain the early maturity and high-yielding essential derivative variety using gamma-ray irradiation. A soybean mutant line from the Kipas Putih cultivar was obtained in the previous study.

The selection in M5 generation **obtained 33 putative mutant** lines, including **6 early maturity mutant lines** (4-14 days earlier than parent), 3 mutant lines that harvested 8 days earlier than a parent, 19 mutant lines with high yields but not early maturity, and **7 putative mutant lines** with high yields but not early maturity and have large seed weight (Nilahayati 2018). Putative mutant lines that are in early maturity are purified for several generations so that they can be continued for the release of new varieties.

The availability of early maturity soybean varieties will overcome the problem of climate change because the use of short-lived varieties will reduce the risk of crop failure due to drought. Furthermore, early maturity soybeans would be more profitable for farmers to alternate crops with rice and avoid water shortages for plants during their growth when planted after rice (Rehajeng and Adie 2013). In addition, early maturity soybeans can provide various advantages, namely reducing pest infestations and increasing the cropping index in a year.

Therefore, early maturity putative mutant lines that have been obtained from the previous study need to be tested in the next generation to determine their genetic stability. The test was carried out by observing the agronomic diversity and yield of putative mutants selected by comparing their characters with the parents and the NILAHAYATI et al. – Several soybean from gamma-rays irradiation in M6 generation 35 comparison varieties. Therefore, a combination of plant breeding and agronomy is needed to improve plant characteristics and test line stability before releasing a variety.

This study used six soybean mutant lines of M6 generation from the Kipas Putih soybean variety mutation with two check varieties, namely Kipas Putih and Anjasmoro. All soybean seeds were obtained from the results of previous generations of research. These lines need to be tested for yield to determine agronomic performance with their parents. MATERIALS AND METHODS The research was conducted in Paloh Lada Village, Dewantara Subdistrict, Aceh Utara District, Indonesia, from July 2021 to October 2021. The plant materials used in this study were eight soybean genotypes, namely Kipas Putih variety (parent variety), Anjasmoro variety (comparison variety), M.1.1.3, M.5.2.1, M.5.2.3, M.1.1.8, M.1.1.9 and M.1.1.17 mutant lines.

The agricultural production facilities used are urea 50 kg ha⁻¹, SP-36 150 kg ha⁻¹, KCl 100 kg ha⁻¹, manure 3 tha⁻¹, Decis 25 EC and Dithane-M45. A non-factorial Randomized Block Design (RBD) was utilized as the experimental design. The planting area is cleaned of weeds that grow in the area. Then made an experimental plot with a size of 2 m x 1.5 m. A drainage ditch was made with a distance between plots and between replications of 50 cm. Seed planting is done by making planting holes in the plot with a depth of 2 cm and a spacing of 40 cm x 20 cm, then inserting 2 seeds per planting hole.

Thinning was done at the age of 2 weeks after planting so that only 1 plant is left per planting hole. Fertilization was carried out according to the recommended dosage of soybean fertilizer. The application of SP-36 and KCl fertilizers and manure was carried out 2 weeks before planting, while urea was applied at the time of planting. Thinning was done at the age of 2 MST. Weeding was done manually by removing weeds from

the plot. Harvesting is done by pulling the stems of the plant by hand. The harvest criteria were that most of the leaves had turned yellow and fell, the skin of the pods was brownish yellow as much as 95% of the experimental plot units.

Observations were made on the variables of plant height, number of branches, flowering age, harvest age, number of pods per plant, seed weight per plant, seed weight per plot and tha^{-1} production. Data analysis was performed by analysis of variance. If significantly different, it continues with the Least Significant Difference (LSD) Test. RESULTS AND DISCUSSION Plant height The results showed that there was a very significant effect on the character of plant height of several M6 genotypes and their two comparison varieties.

The results of further tests on the average plant height of several M6 genotypes and their two comparison varieties can be seen in Figure 1. Figure 1. The effect of several soybean genotypes on plant height characters. Note: The numbers followed by the same letter in the same bar are not significantly different according to the 5% LSD test. The highest plants were found in Kipas Putih variety with a height at harvest reaching 110.50 cm, which was not significantly different from the height of genotype Anjasmoro with a height of 99.66 cm. Then followed by M.5.2.1, M.7.1.17, M. 5.2.3, M.1.1.8 and M.1.1.3 mutant lines. The shortest plant height was found in M.1.1.9

genotype with a height of only 73.94 cm, which was not significantly different from the height of M.1.1.8 and M.1.1.3 mutant lines (Figure 1). In the mutant lines of gamma-ray irradiation, the selection results in the previous generation that were tested still showed a decrease in plant height compared to Kipas Putih. The reduction of plant height due to gamma-ray irradiation treatment on plants has been found by many researchers before. Yadav et al.

(2019) found that maize genotype HQGM-1 irradiated at several doses of irradiation showed a reduction in plant height of 56% at an irradiation dose of 0.5 kGy. (M.T. et al. 2021) also reported the development of semi-dwarf, early maturing and high-yielding mutant of rice cultivar Improved White Ponni with gamma-rays irradiation. In the M6 generation, they found a significant reduction in plant height (up to 40% reduction). Similar results have been reported by (Mudibu et al. 2012) in soybean. In the 0.4 kGy gamma-rays treatment, plant height was seen to be shorter in M1 generation, with a decrease of 25%, 13% and 38% for Kitoko, Vuangi, and TGX814-49D, respectively.

In M2 generation, a 0.4 kGy gamma-ray irradiation dose caused a 14% decrease in plant height in the Kitoko genotype. The plant height reduction in M1 generation was observed for the 0.4 kGy treatment with 25%, 13% and 38% decrease for Kitoko, Vuangi,

and TGX814-49D, respectively. In M2 generation, the 0.4 kGy irradiation significantly reduced plant height in Kitoko (14%). When compared to planting in the previous generation (M5 generation), the plant height of the tested lines ranged from 39-59 cm, which means that it was shorter in height than the planting in this M6 generation.

This is due to different locations and different growing seasons. Plant height is included in the quantitative character which is strongly influenced by the environment. Xue et al. (2019) stated that plant height is an important trait in soybean. The taller plants may give the higher yields but will be more susceptible to lodging. There are many genes involved N U S A N T A R A B I O S C I E N C E 14 (1): 34-39, May 2022 36 together in influencing plant height throughout development. Genetically, plant height is a quantitative trait controlled by multiple genes.

Days of flowering, harvesting age and number of branches The analysis of variance showed a very significant effect on the characteristics of flowering and harvesting ages of several M6 genotypes and their two comparison varieties. However, there was no significant effect on the character of the number of branches of several M6 genotypes and their two comparison varieties. Further tests on the average days of flowering, number of branches and harvesting age, and their two comparison varieties can be seen in Figures 2, 3 and 4. The longest days of flowering was found in M.5.2.3 mutant genotype with a flowering age of 42.77 days after planting (DAP) which was not significantly different from other mutant genotypes and the parent variety. The fastest flowering age was found in Anjasmoro which started flowering at 35.55 DAP.

The research results in this generation showed that there was an unstable appearance on the character of flowering age compared to the planting of these lines in the previous generation. Nilahayati (2018) found that in the M5 generation, the M.1.1.3, M.5.2.1, M.5.2.3 and M.1.1.9 lines were selected based on the criteria for flowering age which were faster than other lines with flowering age at 36-37 DAP. In contrast, M.T. et al. (2021) reported the reduction days to flowering in the M6 generation of Improved White Ponni (IWP) paddy mutants. Days to 50% flowering of WP 5-4 mutant was reduced to 13 days (11.8% reduction from 110 days in the IWP control).

In the other six mutants there was a reduction of up to 11 days in their flowering time. The longest harvesting age was found in genotype Kipas Putih with a harvest age 89.66 DAP, which was significantly different from the harvest age of genotypes M.1.1.17, M.5.2.3 and M.1.1.8. The earliest harvesting age was found in genotype Anjasmoro with a harvest age 82.83 DAP, which was not significantly different from the harvest age of M.1.1.3, M.5.2.1 and M.1.1.9 with harvesting ages 83.89, 83.72 and 83.05 DAP, respectively. When compared with the previous generation, the genotypes of the

M.1.1.3, M.5.2.1 and M.1.1.9 mutant lines showed an early harvesting age.

These mutants can be called solid mutants for harvesting age because they provide stability in the character of early harvest in several generations of planting. The mutant genotypes, namely M.1.1.3, M.5.2.1 and M.1.1.9 had an average harvest age of 83 DAP so they were categorized as early maturity soybeans (78-85 days). The Previous research to obtain early maturity soybeans using gamma-ray irradiation has also been carried out by (Puspitasari et al. 2021) on Argomulyo soybean with a radiation dose of 250 Gray. They get the harvest time of the mutant line was more early (70 days) than the control (>80 days). The GEE-5 mutant line showed the shortest harvest time of 66.79 days.

The mutant line flowering days also showed a faster flowering time (31-32 days), while the control variety was around 35 days. The highest average seed yield was found in mutant lines GBB-2 and GEE-5 compared to control varieties and other mutant lines because this mutant had a higher number of filled pods than other genotypes. However, the seed size of the mutant line was smaller, only 10-11 g compared to the 100 seeds weight of the control varieties, which was 13-16 g. Figure 2. The effect of several soybean genotypes on the characters of days of flowering.

Note: The numbers followed by the same letter in the same bar are not significantly different according to the 5% LSD test Figure 3. The effect of several soybean genotypes on the characters number of branches. Note: The numbers followed by the same letter in the same bar are not significantly different according to the 5% LSD test Figure 4. The effect of several soybean genotypes on the characters of harvesting age. Note: The numbers followed by the same letter in the same bar are not significantly different according to the 5% LSD test NILAHAYATI et al.

– Several soybean from gamma-rays irradiation in M6 generation 37 The preference for early maturity soybeans was higher than late maturity soybeans. This is because it could increase the cropping index. It can also avoid crop failure due to drought stress due to shorter pod filling mass. The length of time for pod filling is the most critical and vulnerable period in the event of a long dry season. Rahajeng and Adie (2013) said that early maturity soybean varieties could be a solution for farmers to deal with climate change.

Early maturing varieties are in great demand because they can provide various benefits, such as eliminating yield losses due to drought and pest infestations and increasing cropping in a year. The number of pod and 100-seeds weight The results of the analysis of variance showed that there was a very significant effect on the character of the number pods and the weight of 100 seeds of several M6 genotypes and their two

comparison varieties. The results of further tests on the average number of pods and weight of 100 seeds on several M6 genotypes and their two comparison varieties can be seen in Figure 5 dan 6.

Figure 5 shows that the highest number of pods was found in Kipas Putih variety, which amounted to 239.89 pods, which was not significantly different from M.1.1.8 mutant line with 217.33 pods. The lowest number of pods was obtained in M.1.1.3 mutant line with only 97.05 pods, which was not significantly different from M.1.1.9, M.1.1.17 lines and the comparison variety Anjasmoro. There was only one putative mutant genotype that had the same number of pods as its parents, namely the M.1.1.8 genotype. There was no genotype whose number of pods exceeded the number of parent pods. Genotype M.1.1.8

in the previous generation test (M5) had the highest number of pods compared to other genotypes, with an average of 500 pods (twice the number of parent pods). In the planting of the M6 generation, there was a decrease in the average number of pithy pods, probably due to the different locations and planting seasons. The largest weight of 100 seeds was found in Anjasmoro variety at 17.34 g/100 seeds (Figure 6). The smallest 100-seeds weight was found in the M.1.1.3 mutant line with 100 seed weight 11.79 g, which was not significantly different from the Kipas Putih variety, the M.1.1.9 and M.1.1.8 mutant lines, which weighted 100 seeds, respectively 11.68 g, 11.82 g and 12.55 g. The mutant line that increased seed size compared to its parents in M6 generation was the M.5.2.3 mutant line, which increased by 2 g/100 seeds.

This mutant line on the previous M5 generation planting showed a small seed size of only 10.21 g/100 seeds. This indicates that differences influence the increased seed size in the planting environment. Soybean seed size was divided into three groups, namely small seed size (<10 g/100 seeds), medium seed size (10-14 g/100 seeds), and seed size >14 g/100 seed as large seed size (Krisnawati and Adie 2015). Based on these criteria, the putative mutant lines in this study had small seed sizes and medium-sized seeds were not different from their parents, Kipas Putih.

At the same time, the comparison variety is Anjasmoro, a large-seed soybean variety with a weight of 100 seeds 17 g, which follows the variety's description. Seed weight/plant, seed weight/plot and production tha-1 The results of the analysis of variance showed that there was a significant to a very significant effect on the character of seed weight/plant, seed weight/plot and production of several M6 genotypes and their two comparison varieties. Further tests on the average seed weight/plant, seed weight/plot and production/ha on several M6 genotypes and their two comparison varieties can be seen in Figures 7, 8 and 9.

Figure 7 dan 9 shows that the highest **seed weight per plant** and production per hectare was found in the Kipas Putih variety with a seed weight of 42.25 g and production of 4.75 tons/h, which was not significantly different from the seed weight of M.1.1.8 mutant line, which was 38.84 g/plant and production of 4.37 tha⁻¹. The lowest seed weight and production were found in M.1.1.3 mutant line with seed weight only 18.25 g and production 2.05 tha⁻¹, which were not significantly different from M.7.1.17 and M.1.1.9. In the M.1.1.9 mutant line, the low **seed weight per plant** was because many pods were contained, but the seeds produced were not good, such as the small and shriveled seeds' shape. Figure 5.

The effect of several soybean genotypes on the characters **number of pods and the weight of 100 seeds**. Note: The **numbers followed by the same letter in the same bar are not significantly different according to the 5% LSD test** Figure 6. The effect of several soybean genotypes on **the weight of 100 seeds** characters. Note: The **numbers followed by the same letter in the same bar are not significantly different according to the 5% LSD test** N U S A N T A R A B I O S C I E N C E 14 (1): 34-39, May 2022 38 Figure 7.

The effect of several soybean genotypes on seed weight/plot characters. Note: The **numbers followed by the same letter in the same bar are not significantly different according to the 5% LSD test** Figure 8. The effect of several soybean genotypes on seed weight/plot characters. Note: The **numbers followed by the same letter in the same bar are not significantly different according to the 5% LSD test** Figure 9. The effect of several soybean genotypes on production tha⁻¹.

Note: The **numbers followed by the same letter in the same bar are not significantly different according to the 5% LSD test** Late-maturing soybeans (Kipas Putih variety) yield more seeds than early-maturing ones because their vegetative growth and reproductive periods are longer. Therefore, it has been difficult to develop an early-maturing line without decreasing seed yield. The present study showed that M.1.1.8 mutant had a significantly earlier maturing time than Kipas Putih (Figure 2), and M.1.1.8 had seed yield similar to Kipas Putih (Figure 9). Therefore, we have successfully found an early-maturing mutant line without decreasing yield.

For measurement data of seed weight/plot (Figure 8), it can be seen that the highest seed weight/plot was found in Kipas Putih variety, namely 489.30 g/plot, which was not significantly different from M.5.2.3, M.5.2.1 and M.1.1.8. On the other hand, the lowest **seed weight per plot** was found in the M.7.1.17 mutant line, which was 303 g/plot. There was a decrease in the seed weight/plots of the M.1.1.8 mutant line because the number of plants/plots that managed to grow in this line was only 50% of the total number of

plants.

From the total 24 plant/plot that managed to grow only 12 plants per plot, resulting in a decrease in seed weight/plot compared to seed/plant weight data. These results are in line with (Asadi and Dewi 2020). They obtained soybean mutants resulting from gamma irradiation combined with crossbreeding. As a result, there were **nine mutant soybean lines** with better performance and 15-26% higher yield than the Panderman variety **and 27-31% higher than the Anjasmoro** variety.

The nine selected lines will be tested for further adaptability in different locations and environments in the next generation. Several mutant soybeans were also obtained through gamma-ray irradiation of the Panderman variety. The results on the M5 generation of 10 tested mutants show two mutant lines, i.e., Kdl3 and Kdl8, that show great performance than the other mutant lines under drought stress conditions. These mutant lines required 30.75 to 32 days to flower and 79.75 to 83.75 days to harvest. They also have relatively short plant heights 28.25 and 23.35 cm, respectively (Yuliasti and Reflinur 2017). In this study, sterile mutants were still found in M.1.1.3 mutant lines population.

There was only one full sterility plant from the entire cultivated population, as shown in Figure 10. When entering the flowering phase, this plant shows the appearance of several flower candidates but in 2 days they fall, so the flower candidates appear can not grow. In addition, this plant shows differences in growth when it reaches harvest age, namely the leaves turn yellow more quickly and over time the leaves wither and fall off.

Several other abnormal characteristics in full sterility plants were the number of branches and the number of leaves was less when compared to other plants. Figure 10. A. Fertile mutant lines, B. Sterile mutant lines A B NILAHAYATI et al. – Several soybean from gamma-rays irradiation in M6 generation 39 In a previous study, Nilahayati et al. (2016) also reported that full sterility plants showed normal vegetative growth but failed to produce flowers. Kumar and Rai (2006) investigated the **study on soybean using** gamma- rays irradiation and found some sterile mutants. The mutant cytology test showed the occurrence of desynapsis in the cells.

In sterile plants, only slightly bivalent while the univalent frequency is very high. This causes a very high percentage of pollen sterility. The later stages of meiosis are also severely disrupted. Plants were identified as male sterile plants because only a few pods were formed. They concluded that gamma-rays may have acted on some of the genes responsible for the formation of synapses and chiasms at the stage of gamete cell division. In conclusion, the genotype treatment affected all observed characters except

for the number of branches and the number of empty pods. The putative mutant line with high yield and equivalent to its parents is M.1.1.8 line with production 4.37 tons/ha.

The putative mutant lines with early maturity were M.1.1.9, M.5.2.1 and M.1.1.3 lines with a harvesting age of 83 DAP. ACKNOWLEDGEMENTS This research was funded by Universitas Malikussaleh in accordance with Decree Number 530/UN45/KPT/2021, July 12, 2021 and Agreement/Contract Number 95/PPK- 2/SPK-JL/2021. REFERENCES Asadi, Dewi N. 2020. Agronomic performance of soybean mutant lines based on preliminary and advance yield at dryland area. The 1st International Conference on Genetic Resources and Biotechnology. IOP Conf Ser Earth Environ Sci 482 (2020): 012018. DOI: 10.1088/1755-1315/482/1/012018.

Harsanti L, Parno, Tarmizi. 2020. The effect of gamma Co 60 radiation on soybean (*Glycine max* (L.) Merr) shade tolerants. IOP Conf Ser Earth Environ Sci 518 (2020): 012055. DOI: 10.1088/1755- 1315/518/1/012055. Harsono A, Pratiwi H. 2017. The increase of ultisol productivity based on intercropping cassava with peanut and soybean. Nusantara Biosci 9: 157-163. DOI: 10.13057/nusbiosci/n090209. Hong MJ, Jang YE, Kim DG, Kim JM, Lee MK, Kim JB, Eom SH, Ha BK, Lyu J, Kwon SJ. 2019. Selection of mutants with high linolenic acid contents and characterization of fatty acid desaturase 2 and 3 genes during seed development in soybean (*Glycine max*). J Sci Food Agric 99 (12): 5384-5391. DOI: 10.1002/jsfa.9798.

Hussain SZ, Akram G, Shabbir, Ahmad Q. 2020. Response of soybean genotypes to different levels of mutagens for yield related attributes. Biomed J Sci Tech Res 25 (1): 18757-18764. DOI: 10.26717/BJSTR.2020.25.004135. Krisnawati A, Adie AA. 2015. Selection of soybean genotypes by seed size and its prospects for industrial raw material in Indonesia. Proc Food Sci 3: 355-363. DOI: 10.1016/j.profoo.2015.01.039. Kumar G, Rai P. 2006. Induced desynaptic male sterile lines in soybean. Cytologia 71 (4): 337-343. DOI: 10.1508/cytologia.71.337. Lestari SAD, Sutrisno, Kuntiyastuti H. 2017. Short Communication: Effect of organic and inorganic fertilizers application on the early-medium maturing soybean yield. Nusantara Biosci 10: 1-5.

DOI: 10.13057/nusbiosci/n100101. Malek MA, Rafii MY, Afroz SS, Nath UK, Mondal MMA. 2014. Morphological characterization and assessment of genetic variability, character association, and divergence in soybean mutants. Sci World J 2014 (1): 968796. DOI: 10.1155/2014/968796. M.T. A-P-L, Ramchander S, Kumar KK, Muthamilarasan M, Pillai MA. 2021. Assessment of efficacy of mutagenesis of gamma-irradiation in plant height and days to maturity through expression analysis in rice. PLoS One 16 (1): e0245603. DOI: 10.1371/journal. pone.0245603. Mawarni L, Nisa TC, Napitupulu JA, Karyudi. 2019. Determination of leaf status of soybean varieties on shading. Biodiversitas

20: 615-620. DOI: 10.13057/biodiv/d200243. Mudibu, J.

Nkongolo CKK, Kalonji-Mbuyi A, Kizungu RU. 2012. Effect of gamma radiation on morpho-agronomic characteristic of soybeans (*Glycine max L.*). *Am J Plant Sci* 3: 331-337. DOI: 10.4236/ajps.2012.33039. Nobre DAC, Macedo WR, Sedyama CS, Piovesan N. 2019. Gamma radiation to produce soybean mutants for better plant performance and chemical composition of seeds. *Emir J Food Agric* 31 (7): 511- 519. DOI: 10.9755/ejfa.2019.v31.i7.1983. Nilahayati, Rosmayati, Hanafiah DS, Harahap F. 2016. Gamma irradiation induced chlorophyll and morphological mutation in Kipas Putih soybean. *Intl J Sci Basic Appl Res* 30 (3): 74-79. Nilahayati, 2018. The Improvement of Agronomic Character of Kipas Putih Soybean through Gamma-ray Irradiation. [Dissertation].

Universitas Sumatera Utara, Medan. [Indonesian]. Nilahayati, Rosmayati, Hanafiah DS, Harahap F. 2019. The genotype selection of M3 generation of Kipas Putih soybean with gamma-rays irradiation on agronomic characters, early maturity and high yielding mutants. *Bulg J Agric Sci* 25 (1): 97-102. Puspitasari W, Arwin, Yuliasti. 2021. Agronomic characters of early maturity soybean mutant lines. *AIP Conf Proc* 2381: 020068. DOI: 10.1063/5.0066925. Rahajeng W, Adie MM. 2013. Varietas kedelai umur genjah. *Buletin Palawija* 26: 91-100. DOI: 10.21082/bulpa.v0n26.2013.p91-100.

[Indonesian] Toyip, Ghulamahdi M, Sopandie D, Aziz SA, Sutandi A, Purwanto MYJ. 2019. Physiological responses of four soybean varieties and their effect to the yield in several saturated soil culture modification. *Biodiversitas* 20: 2266-2272. DOI: 10.13057/biodiv/d200822. Xue H, Tian X, Zhang K, Li W, Qi Z, Fang Y, Li X, Wang Y, Song J, Li WX, Ning H. 2019. Mapping developmental QTL for plant height in soybean [*Glycine max (L.) Merr.*] using a four-way recombinant inbred line population. *PLoS One* 14 (11): e0224897. DOI: 10.1371/journal.pone.0224897. Yadav A, Singh B, Singh SD. 2019. Impact of gamma irradiation on growth, yield and physiological attributed of maize. *Indian J Exp Biol* 57: 116-122. Yuliasti, Reflinur. 2017.

Field performance of five soybean mutants under drought stress conditions and molecular analysis using SSR markers. *Atom Indones* 43 (2): 103-109. DOI: 10.17146/aij.2017.685. Yoel A, Rachmadi M. Morphological characteristics and yield performance of mutant soybean [*Glycine max (L.) Merr.*] M6 genotypes in Jatinangor, Sumedang, West Java. *Agric Sci* 5 (3): 117- 122. DOI: 10.22146/ipas.54104.

INTERNET SOURCES:

<1% - www.academia.edu > es > 41843139

<1% - pdfs.semanticscholar.org › 416a › 2a928060576b23aa9b
1% - jthort.org › index › jthort
<1% - ui.adsabs.harvard.edu › abs › 2018E&ES
<1% - www.embibe.com › exams › plant-breeding
<1% - www.cambridge.org › core › journals
<1% - repositori.usu.ac.id › handle › 123456789
<1% - www.ukm.my › jsm › pdf_files
<1% - sabraojournal.org › wp-content › uploads
<1% - www.ncbi.nlm.nih.gov › pmc › articles
<1% - pdfs.semanticscholar.org › 5cbd › 7f386c1e
<1% - www.reddit.com › r › PlantedTank
<1% - www.researchgate.net › profile › Wayan-Wangiyana
<1% - www.researchgate.net › figure › Analysis-of-variance
<1% - link.springer.com › referenceworkentry › 10
2% - www.researchgate.net › figure › Numbers-followed-by
<1% - nopr.niscair.res.in › bitstream › 123456789/45754/1
<1% - www.semanticscholar.org › paper › Assessment-of
<1% - link.springer.com › article › 10
<1% - file.scirp.org › Html › 4-2600283_17913
<1% - quizlet.com › 69795586 › bio-111-unit-1-intro-to
<1% - researchgate.net › figure › Mean-days-to-flowering
<1% - www.researchgate.net › figure › Plant-height-at
<1% - www.researchgate.net › figure › Condition-of-the
<1% - www.researchgate.net › figure › Mean-variability-and
<1% - researchgate.net › figure › The-number-of-pods-seeds
<1% - www.assignmentexpert.com › homework-answers
<1% - www.researchgate.net › figure › Number-of-pods-per
<1% - www.sciencedirect.com › science › article
<1% - pubmed.ncbi.nlm.nih.gov › 30369935
<1% - www.researchgate.net › figure › Interaction-effect
<1% - www.researchgate.net › figure › Seed-weight-per-plot
<1% - repository.pertanian.go.id › 12054 › soybean2020
<1% - www.researchgate.net › publication › 247836752
<1% - www.researchgate.net › publication › 341738777
<1% - www.researchgate.net › publication › 346001557_The
<1% - www.semanticscholar.org › paper › The-increase-of
1% - www.ncbi.nlm.nih.gov › pubmed › 31077382
<1% - biomedres.us › pdfs › BJSTR
<1% - www.litbang.pertanian.go.id › artikel › 345
2% - smujo.id › biodiv › article

1% - journals.plos.org › plosone › article
<1% - www.scirp.org › (S(i43dyn45teexjx455qlt3d2q
<1% - www.semanticscholar.org › paper › Gamma-radiation-to
<1% - www.latlong.net › place › universitas-sumatera-utara
<1% - www.researchgate.net › publication › 332092976_The
<1% - www.semanticscholar.org › paper › Agronomic
<1% - pubmed.ncbi.nlm.nih.gov › 27918577
<1% - www.semanticscholar.org › paper › Field-Performance
<1% - doaj.org › article › 6a7a8ecf0a224f62b16bcbfaf8ec5ab6